

For Acne Vulgaris, A Single Agent That's Easy to Apply

EVOCLIN[®] (clindamycin phosphate) Foam, 1%




evoclin[®]
(clindamycin
phosphate) Foam, 1%

Please see Important Safety Information on back page
and accompanying Full Prescribing Information.

With its foam delivery system, EVOCLIN® (clindamycin phosphate) Foam, 1% provides easy, spreadable coverage of hair-bearing and hairless body parts, with minimal residue. The foam dissipates quickly, does not drip, is nonstaining, nonbleaching, and fragrance free.

EVOCLIN Foam is indicated for topical application in the treatment of acne vulgaris in patients 12 years and older. It is for topical use only and not for oral, ophthalmic, or intravaginal use. Avoid contact with eyes, mouth, lips, other mucous membranes, or areas of broken skin. If contact occurs, rinse thoroughly with water. In view of the potential for diarrhea, bloody diarrhea, and pseudomembranous colitis, the physician should consider whether other agents are more appropriate.

The foam is applied once daily to affected areas after the skin is washed with mild soap and allowed to fully dry. The entire affected area should be covered with foam. If there has been no improvement after 6 to 8 weeks or if the condition becomes worse, treatment should be discontinued.



DISCOUNTS FOR YOUR PATIENTS

PRESTIUM PRESCRIPTION SAVINGS PROGRAM

With Prestium Pharma's Prescription Savings Program, your patients can save up to \$45 off out-of-pocket costs each time their pharmacist fills or refills the prescription. **Ask for a supply of Prescription Savings Cards (see sample below) to provide your patients with their script.**

The cards are valid at participating pharmacies for one use per medicine per month. Not valid for patients under Medicaid, Medicare (including Medicare Part D), or similar state or federal programs. The card is not valid for residents of Massachusetts unless the patient is paying the full cost of the prescription.

INSTANT SAVINGS CARD
Pay the initial \$15 and receive up to \$45 off your out-of-pocket expenses.

DenaVir[®] (periclovir cream, 1%)	ELIMITE[™] CREAM (Pimecrolimus) 5%	evoclin (indinavir prochlorate) foam, 1%	extina[®] (ketotifen) foam, 1%
Luciq[®] (bimatoprost) foam, 0.12%	Olux[®] (betamethasone propionate) foam, 0.05%	Olux-E (betamethasone propionate) foam, 0.05%	Vision (0.2% w/v benzalkonium chloride, 1% w/v polyvinyl alcohol, 0.2% w/v polyvinylpyrrolidone) solution

BIN: 610020 GROUP: 99992198 ID: XXXXXXXXXXXX
Please see back of card for details.



How the Prestium Pharma Combo Savings Card Works for Your Patients:

- Pay the initial \$15, you could receive up to \$45 off your co-pay or out-of-pocket expenses.
- You may pay more than \$15 if your co-pay exceeds \$60; if your insurance does not cover Denavir, Elimite, Evoclin, Extina, Luxiq, Olux, Olux-E or Vusion; or if you are a cash payer.
- Go to www.PrestiumPharma.com for more information.
- For assistance with the Prestium Pharma Combo Savings Card, contact **1-855-820-3232**.
- After each transaction, keep this card for up to 12 total uses.

**APPLY FOR YOUR
PRESCRIPTION CARD TODAY!**



IMPORTANT SAFETY INFORMATION

Please see accompanying Full Prescribing Information

- EVOCLIN® (clindamycin phosphate) Foam, 1% is contraindicated in individuals with a history of hypersensitivity to preparations containing clindamycin or lincomycin, or a history of regional enteritis, ulcerative colitis, or antibiotic-associated colitis, including pseudomembranous colitis.
- Prescribe with caution in atopic individuals.
- Concomitant topical acne therapy should be used with caution since a possible cumulative irritancy may occur, especially with the use of peeling, desquamating, or abrasive agents. If irritation or dermatitis occurs, EVOCLIN Foam should be discontinued.
- Diarrhea, bloody diarrhea, and pseudomembranous colitis have been reported with systemic and rarely with topical clindamycin. Discontinuation is recommended if significant diarrhea develops.
- In clinical trials, the most common adverse events were headache (3%) and application-site reactions, including burning (6%), itching (1%), and dryness (1%).
- The contents of EVOCLIN Foam are flammable; patients should avoid fire, flame, and/or smoking during and immediately following application.



evoclin[®]

(clindamycin
phosphate) Foam, 1%

Prestium[®]
Pharma

prestiumpharma.com

Prest-Evoclin-001